

Stability of the hydrogen atom of classical electrodynamics

Jayme De Luca*

Universidade Federal de São Carlos,

Departamento de Física

Rodovia Washington Luis, km 235

Caixa Postal 676, São Carlos, São Paulo 13565-905

(Dated: February 2, 2008)

We study the stability of the circular orbits of the hydrogen atom in a version of classical electrodynamics that was historically overlooked. We introduce the concept of resonant dissipation, i.e. a motion that radiates the center-of-mass energy while the interparticle distance performs bounded oscillations about a metastable orbit. The stability mechanism is established by the existence of a quartic resonant constant generated by the stiff eigenvalues of the linear stability problem. This constant bounds the particles together during the radiative recoil. The condition of resonant dissipation predicts angular momenta for the metastable orbits in reasonable agreement with the Bohr atom. The principal result is that the emission lines agree with the predictions of quantum electrodynamics (QED) with 1 percent average error even up to the 40th line. Our angular momenta depend logarithmically on the mass of the heavy body, such that the deuterium and the muonium atoms have essentially the same angular momenta, in agreement with QED. Analogously to QED, our stability analysis naturally uses the eigenvalues of an infinite-dimensional linear operator; the infinite-dimensionality is brought in by the delay of the electromagnetic interaction.

According to classical electrodynamics, a quasi-circular orbit of the hydrogen atom radiates energy. We show that this complex Poincaré-invariant two-body motion can damp the center-of-mass energy only, while the interparticle distance performs bounded oscillations about some circular orbits, a phenomenon henceforth called resonant dissipation. This dynamics emits radiation of a sharp frequency while the center-of-mass energy decreases to compensate for the energy damping, during the life-time of the emission process, i.e. a collective radiative recoil. The stability mechanism is established by the existence of a resonant constant that keeps the interparticle distance bounded during the radiative recoil and deformation. This resonant normal form exists because of a resonance involving the stiff modes of the linearized delay equations of motion, i.e., the linear normal modes with eigenvalues of a very large imaginary magnitude [1]. The results of this simple stability analysis contain no adjustable parameter, satisfy Maxwell's equations and predict the magnitudes of the Bohr atom[2] with accuracy and qualitative detail. We calculate every circular emission line of hydrogen with a 1 percent error. In this paper we also introduce an electromagnetic setting slightly different from the most popular form of electrodynamics with self-interaction and retarded fields. This most popular form is henceforth called the dissipative-retarded setting (DR). Here we take the equations of motion derived from Fokker's Lagrangian of the action-at-a-distance electrodynamics [1, 3, 4], with advanced and retarded interactions, and we add the Lorentz-Dirac self-interaction force[5] on each particle. This new electromagnetic setting is henceforth called the dissipative-Fokker model (DF). Both the DR and the DF settings satisfy Maxwell's equations everywhere and define covariant dynamical systems for the electrodynamics of classi-

cal point charges. We experimented with the stability analysis of the several possibilities within classical electrodynamics, and DF is the most transparent setting for the resonant dissipation phenomenon introduced here. Our results advocate for the interest of DF; we predict the Bohr atom with great precision and qualitative detail. It is popular to use the word radiation as equivalent to dissipation, because DR electrodynamics is absolutely dissipative. There is nevertheless a subtle difference from Poyting's theorem inside DF to Poyting's theorem inside DR; the energy flux is not necessarily dissipative in DF (because we have advanced and retarded Liénard-Wiechert far-fields). This allows the atom to receive energy from other atoms of the universe via the effective half/advanced plus half/retarded far-fields, a thermalization mechanism that is absent in DR.

To understand how our dynamics is a solution of the equations of motion of classical electrodynamics, one needs to go beyond the simplifications suggested by the Galilei-invariant Coulomb problem. In quantum electrodynamics (QED) the semiclassical Bohr orbits of hydrogen correspond to excited quantum states that decay to the ground state in a life-time of about 10^6 turns[2]. The analog of this decay process is troublesome in DR and subtle in DF. Disregarding the singular coupling to the center-of-mass coordinate is a naive use of the equations of motion of DR, which predicts that the radiated energy comes out of the electrostatic potential energy. In this process the interparticle distance decreases while the rotation frequency increases by a factor of two during the life-time, making the emission of a sharp line impossible. It is very instructive to estimate this energy dissipation along hypothetical circular orbits directly from the Lorentz-Dirac self-interaction force[5]. The complexity disregarded by this simplifica-

tion stems from three main sources: (i) The group theoretical complexity of Poincaré-invariant two-body dynamics; the two no-interaction theorems[6] limit Hamiltonian and local Lagrangian descriptions of two-body dynamics to straight-line free motions only. These negative theorems are sound warning that the Galilei-invariant Coulombian dynamics, with trivial and separable center-of-mass motion, is a bad approximation to relativistic two-body dynamics. (ii) The delay equations of motion bring in an infinite-dimensional dynamical system and needs an initial function as the initial condition; addition of the non-runaway condition leads to a well-posed mathematical problem [7]. The stability analysis of such dynamics reveals stiff eigenvalues with an arbitrarily large imaginary part, as long as integer multiples of π (an arbitrary integer number appears naturally) [1]. (iii) The generic solution to a stiff differential equation is not slowly varying everywhere and typically it has jumps; see for example the equations of Lienard type in Ref.[8]. Another example is the small parameter multiplying the highest derivative of the Lorentz-Dirac self-interaction force[5]. After Dirac's derivation of a finite force of self-interaction within classical electrodynamics[5], Eliezer considered a model two-body motion with Coulomb interaction plus self-interaction[9]. A simple physical consequence transpired, that with self-interaction the electron cannot fall into the proton even along unidimensional collision orbits. This surprising result is a warning that self-interaction is a singular perturbation over the Coulombian dynamics and also suggests that colinear orbits are the natural attractors of the dissipative dynamics (a ground state with zero angular momentum). Along such orbits the heavy particle (the proton) moves in a non-Coulombian way and the self-interaction provides a repulsive mechanism that avoids a collision. Disregard of the stiff nature of the equations of classical electrodynamics has produced several conundrums, and the present work is an attempt to recognize the correct qualitative behavior of the solutions of classical electrodynamics.

In Ref. [10] we studied the classical helium atom in the DR setting. Because the very special circular orbits of helium do not radiate in dipole inside DR, the physics of the emission of a sharp line is easy to understand even in the DR setting[10]. There is already an interesting resonant structure for these circular orbits of helium within the Darwin approximation (a low-velocity approximation to Fokker's Lagrangian) [10, 11]. The predictions of the Darwin approximation were tested numerically in Ref. [12], finding non-ionizing states in the atomic magnitude, as predicted in Ref. [10]. The generalization to the circular orbits of the hydrogen atom in the DR setting is troublesome; circular orbits of hydrogen are stable within the Darwin approximation and radiate in dipole, such that a sharp emission line seems like a conundrum for classical electrodynamics in the DR setting. In this work we come to grips with the physics of the hydrogen atom by

changing to DF *and* by understanding that the correct description comes only after the inclusion of the nuclear motion (i.e. a two-body system with delay and dissipation). The resonant dissipation radiates mostly the center-of-mass energy while a slow deformation of circularity and a collective recoil take place. After a possibly long life-time, this resonant dissipation produces enough deformation to destabilize the orbit, i.e. only metastability is possible (this is also what QED says about circular orbits).

Our perturbation theory starts from the unperturbed circular orbit of the isolated electromagnetic two-body problem derived from Fokker's Lagrangian, as explained in Refs.[1, 3, 4]. A motion of the particles in concentric circles with a constant angular speed and along a diameter is a solution of these equations of motion. This is because the symmetric contributions from future and past generate a resulting force that is normal to the velocity of each particle. We henceforth call this orbit the circular orbit[1, 3, 4]. The quantities for this uniform circular motion are given in [3]. Our electron has mass m_1 and travels a circle of radius r_1 while the proton has mass m_2 and travels a circle of radius r_2 . We henceforth use a unit system where the electronic charge is $e = -1$ and the speed of light is $c = 1$. The circular orbit is defined by the retardation angle θ that one particle turns while the light emanating from the other particle reaches it (one light-cone distance away). The frequency Ω of angular rotation is given, to lowest order in θ , by Kepler's law

$$\Omega = \mu\theta^3 + \dots \quad (1)$$

where $\mu \equiv m_1 m_2 / (m_1 + m_2)$ is the reduced mass. For shallow-energy atomic orbits, the angular momentum l_z is given to first order by $l_z = \theta^{-1}$ [3], which is of the order of the inverse of the fine structure constant, $\alpha^{-1} = 137.036$. To simplify the stability analysis, we transform the particle coordinates from Cartesian to gyroscopic coordinates

$$\begin{aligned} x_k + iy_k &\equiv \exp(i\Omega t)[d_k + \eta_k], \\ x_k - iy_k &\equiv \exp(-i\Omega t)[d_k + \xi_k], \end{aligned} \quad (2)$$

where η_k and ξ_k are complex numbers defining the perturbation of the circularity. The coordinates of the electron are defined by Eq. (2) with $k = 1$ and $d_1 = r_1$ while the coordinates of the proton are defined by Eq. (2) with $k = 2$ and $d_2 = -r_2$ (at the same time in the inertial frame the particles are in diametrically opposed positions along the unperturbed orbit [3]). We also need here the stability along the perpendicular z direction, a linear problem that is uncoupled from the xy stability and easier to work out. The linear stability analysis proceeds like in Ref. [1], by substituting the circular orbit plus a perturbation, Eq. (2), into Fokker's Lagrangian, expanding

this Lagrangian to quadratic order and taking the Euler-Lagrange equations to linear order[16]. Because we have a delay system at hand, the normal mode condition (the characteristic equation) has infinitely many roots. This characteristic equation involves hyperbolic functions that can become arbitrarily large and are insensitive to an arbitrarily large imaginary part of the eigenvalue [1]. A standard technique of delay analysis is to keep only the largest powers of the eigenvalue plus the hyperbolic terms in the characteristic equation[14]. This large-imaginary-part stiff limit is performed in Ref. [1] for the equal-mass two-body system (see Eq. (15) of Ref. [1]). Here we generalize this large-imaginary-part limit to the arbitrary-mass case and we computed the characteristic equation with a symbolic algebra software. We define the normal mode eigenvalue by $\lambda\Omega/\theta$, i.e. every coordinate perturbation oscillates in time as $\exp(\lambda\Omega t/\theta)$ (λ is an arbitrary complex number). The limiting form of the characteristic equation for the isolated different-mass case is

$$\left(\frac{\mu\theta^4}{M}\right) \cosh^2(\lambda) = 1, \quad (3)$$

where μ is the reduced mass and $M \equiv m_1 + m_2$. Both the planar and the perpendicular linearized equations have the same limiting characteristic Eq. (3) along circular orbits. For hydrogen (μ/M) is a small factor of about $(1/1824)$. It is important to understand the structure of the roots of Eq. (3) in the complex λ plane, specially for θ of the order of the fine structure constant. It is easy to see that the very small parameter $\frac{\mu\theta^4}{4M} \sim 10^{-13}$ multiplying the hyperbolic cosine on the left-hand side of Eq.(3) determines that $\sigma \equiv |\text{Re}(\lambda)| \simeq \ln(\sqrt{\frac{4M}{\mu\theta^4}})$. For the first 13 excited states of hydrogen this σ is in the interval $14.2 < |\sigma| < 18.2$. The imaginary part of λ can be an arbitrarily large multiple of π , such that the general solution to Eq. (3) is

$$\lambda = \pm(\sigma + i\pi q), \quad (4)$$

where q is an arbitrary integer. The plus or minus sign of Eq. (4) is related to the time-reversibility of the isolated two-body system, a symmetry that is broken by radiation.

Next we include the dissipation of the DF electrodynamics, i.e. the Lorentz-Dirac self-interaction, a calculation performed by adding the self-interaction force to the Lagrangian equations of motion of the isolated system. Here we give only the limiting form of the characteristic equation

$$\left(1 - \frac{g}{\lambda^2}\right) \left(\frac{\mu\theta^4}{M}\right) \cosh^2(\lambda) = 1 - \frac{2}{3}\theta^2\lambda + \frac{1}{\lambda} \left(\frac{\mu\theta^4}{M}\right) \sinh(2\lambda) + \dots \quad (5)$$

where $g = -7$ for the planar stability and $g = 1$ for the perpendicular stability. We give the terms up to

$O(1/\lambda^2)$ because these are essential for the mathematical phenomenon presented. The coefficient g of the term of order $1/\lambda^2$ is the only difference between the planar and the perpendicular characteristic equations up to $O(1/\lambda^2)$. Further terms of size $1/\lambda^3$ and $\theta^4\lambda^2$ produce only small quantitative changes for σ in the atomic range and will be given elsewhere. The linear term on the right-hand side of Eq. (5) with the $2/3$ coefficient is due to the self-interaction force. This dissipative term breaks the time-reversal symmetry of Eq. (3) and the roots of Eq. (5) no longer come in plus or minus pairs. In the following we implement the necessary condition for a convex analytic function of the interparticle distance to exist.

In the same way already used in Ref.[10], this condition should involve the perpendicular direction z as well. This is because if the atom is to recoil like a rigid body while it radiates, one expects the orbital plane to nutate and precess, typical gyroscopic motions of a rigid body. To introduce the main idea simply, we assume that the non-runaway condition restricts the physics to a finite number of linear modes. We take a perpendicular normal mode and a planar normal mode of Eq. (5), of eigenvalues λ_z and λ_{xy} respectively. Suppose we can find these modes such that $\lambda_z + \lambda_{xy} + \lambda_z^* + \lambda_{xy}^* = 0$, which is the necessary condition for a quartic normal form. i.e.

$$\text{Re}(\lambda_z + \lambda_{xy}) = 0. \quad (6)$$

The coordinate of the planar normal mode is a linear combination of the four $\eta\xi$ gyroscopic coordinates: $u \equiv a_{1k}\eta_k + b_{1k}\xi_k$, while the coordinate of the perpendicular z normal mode is $Z \equiv b_1z_1 + b_2z_2$. Because Fokker's Lagrangian is real, λ_z^* and λ_{xy}^* are also solutions to Eq. (5) with complex conjugate normal mode coordinates. Using these normal mode coordinates and Eq. (6), one can show that the following quartic form is a constant of motion up to higher order terms[11]:

$$C \equiv |u|^2|Z|^2 + \dots \quad (7)$$

Notice from Eq. (2) that $|u|^2$ is a quadratic function of the planar coordinates, such that the resonant constant of Eq. (7) limits the excursion along the z direction times the planar distance to the circular orbit, essentially locking the dynamics to the circular orbit. This necessary condition and the continuation of the leading term (7) to an asymptotic series is discussed in Ref. [11]. The lesson of the above construction is that if the interparticle distance is to be bound by a resonant constant while the dissipation goes on, then we must be able to find a pair of roots to Eq. (5) such that Eq. (6) holds. This implies that Eq. (5) has a pair of almost-symmetric roots of the form

$$\begin{aligned} \lambda_{xy} &\equiv (\sigma + \pi qi + i\epsilon_1), \\ \lambda_z &\equiv -(\sigma + \pi qi + i\epsilon_2), \end{aligned} \quad (8)$$

where ϵ_1 and ϵ_2 can be assumed real if $\lambda_z + \lambda_{xy}$ is purely imaginary by absorbing the real part of the perturbation in the definition of σ . The above-defined root-searching problem is well posed and for each integer q conditions (5) and (8) determine θ as a function of q , i.e., θ must be quantized! An asymptotic solution to condition (8) can be obtained by expanding Eqs. (5) up to quadratic order in ϵ_1 and ϵ_2 while treating σ as an approximate constant. This approximation determines the following discrete values for θ

$$\theta^2 = \frac{6(\pi^2 q^2 - \sigma^2)}{\sigma(\pi^2 q^2 + \sigma^2)^2}, \quad (9)$$

and

$$(\epsilon_1 - \epsilon_2) = \frac{4\pi q(3\sigma^2 - \pi^2 q^2)}{\sigma(\sigma^2 + \pi^2 q^2)^2}. \quad (10)$$

According to QED, circular Bohr orbits have maximal angular momentum and a radiative selection rule ($\Delta l = \pm \hbar$) restricts the decay from level $k+1$ to level k only, i.e. circular orbits emit the first line of each spectroscopic series (Lyman, Balmer, Ritz-Paschen, Brackett, etc...), henceforth called the QED circular line. We have solved Eqs. (8) and (5) with a Newton method in the complex λ plane. Every angular momentum $l_z = \theta^{-1}$ determined by Eq. (6) has a value in the correct atomic magnitude, but the subset of Table 1 has frequencies w_{DF} surprisingly close to the QED lines. These lines are when the integer q is approximately equal to an integer multiple of the integer part of 2σ . The reason for such selection rule is that from the resonances satisfying the necessary condition, only some have $|u|^2$ depending on the translation-invariant quantities $(\xi_1 - \xi_2)$ and $(\eta_1 - \eta_2)$ to allow a recoiling translation[16]. In our description the emission mechanism is at a frequency equal to the orbital frequency Ω corrected by the frequency of the complex amplitude uZ defined above Eq. (7), as we explain below. The numerically calculated angular momenta $l_z = \theta^{-1}$ for this select subset are given in Table 1, along with the orbital frequency in atomic units $(137^3\Omega)/\mu = (137\theta)^3$, the QED first frequency of the series in atomic units $w_{QED} \equiv \frac{1}{2}(\frac{1}{k^2} - \frac{1}{(k+1)^2})$, and the frequency predicted by the dissipative Fokker model $w_{DF} \equiv (137\theta)^3 + 137^3\theta^2(\epsilon_1 - \epsilon_2)$. We list only the first 13 lines, which are the experimentally observable, but we tested the agreement of the numerical calculations of the Newton method with up to the 40th circular line predicted by QED. Beyond that, the asymptotic formula (9) shows that the agreement is essentially for any integer k because substitution of $q = [2\sigma]k$ into Eq. (9) yields

$$\theta^{-1} = \sqrt{\frac{2\pi^2}{3}}\sigma^{3/2}k \sim 137.9k, \quad (11)$$

to be compared with the 137.036 of QED. The agreement for any integer k suggests that Eq. (5) is equivalent to

Schroedinger's equation (both are eigenvalue problems for an infinite-dimensional linear operator, and linear operators with the same spectrum are equivalent).

$l_z = \theta^{-1}$	$(137\theta)^3$	w_{QED}	w_{DF}
161.21	6.137×10^{-1}	3.750×10^{-1}	3.688×10^{-1}
282.88	1.136×10^{-1}	6.944×10^{-2}	6.806×10^{-2}
397.40	4.097×10^{-2}	2.430×10^{-2}	2.470×10^{-2}
519.59	1.833×10^{-2}	1.125×10^{-2}	1.113×10^{-2}
637.79	9.911×10^{-3}	6.111×10^{-3}	6.054×10^{-3}
751.48	6.059×10^{-3}	3.685×10^{-3}	3.719×10^{-3}
871.84	3.880×10^{-3}	2.391×10^{-3}	2.392×10^{-3}
987.25	2.672×10^{-3}	1.640×10^{-3}	1.654×10^{-3}
1109.18	1.188×10^{-3}	1.173×10^{-3}	1.170×10^{-3}
1225.92	1.139×10^{-3}	8.678×10^{-4}	8.694×10^{-4}
1343.20	1.061×10^{-3}	6.600×10^{-4}	6.628×10^{-4}
1466.89	8.146×10^{-4}	5.136×10^{-4}	5.102×10^{-4}
1585.51	6.455×10^{-4}	4.076×10^{-4}	4.052×10^{-4}

Table 1: Numerically calculated angular momenta $l_z = \theta^{-1}$ in units of e^2/c , orbital frequencies in atomic units $(137\theta)^3$, circular lines of QED in atomic units $w_{QED} \equiv \frac{1}{2}(\frac{1}{k^2} - \frac{1}{(k+1)^2})$ and the emission frequencies of the DF model in atomic units $w_{DF} \equiv (137\theta)^3 + 137^3\theta^2(\epsilon_1 - \epsilon_2)$.

Last we calculate the frequency of the emission line. The treatment is a consequence of the above defined DF version of electrodynamics and differs from the usual electromagnetic model (DR) in non-trivial ways. We have consistently used the relativistic action-at-a-distance electrodynamics as a many-body electromagnetic theory, as first suggested in Ref. [13]. This no-field theory is based on a parametrization-invariant action involving two-body interactions only, without the mediation by fields. The isolated electromagnetic two-body problem, away from the other charges of the universe, is a time-reversible dynamical system defined by Fokker's action

$$S_F = - \int m_1 ds_1 - \int m_2 ds_2 + \int \int \delta(\|x_1 - x_2\|^2) \dot{x}_1 \cdot \dot{x}_2 ds_1 ds_2, \quad (12)$$

where $x_i(s_i)$ represents the four-position of particle $i = 1, 2$ parametrized by its arc-length s_i , double bars stand for the four-vector modulus $\|x_1 - x_2\|^2 \equiv (x_1 - x_2) \cdot (x_1 - x_2)$, and the dot indicates the Minkowski scalar product of four-vectors with the metric tensor $g_{\mu\nu}$ ($g_{00} = 1, g_{11} = g_{22} = g_{33} = -1$). The Euler-Lagrange equations of Lagrangian (12) were already used to derive our Eq. (3). Surprisingly, the solution to the unusual equations of motion of Lagrangian (12) are still determined by initial position and velocity, as proved in Ref.([7]) for the

non-runaway solutions. Therefore we are dealing with a perfectly well-posed and causal system dressed in an unfamiliar form, and we invite the reader to consult Ref. [7]. In the many-body action-at-a-distance electrodynamics one needs a prescription to describe the incoherent interaction with the universe, because including every single particle of the universe in a pairwise many-body Lagrangian like Eq. (12) is impractical (even though it is precisely what we want to describe physically). This prescription for the dissipative interaction with the universe must be a covariant prescription that satisfies Maxwell's equations. For example, the Wheeler-Feynman prescription of Ref. [13] leads to the usual electromagnetic DR model. The prescription of the DF model is to add the Lorentz-Dirac self-interaction directly to the Euler-Lagrange equations of action (12).

In the DF model the interaction with a distant particle involves half/retarded plus half/advanced Liénard-Wiechert interactions, as derived from a pairwise Lagrangian of type (12). We shall show that this does not preclude radiation of energy. Once DF satisfies Maxwell's equations everywhere, Poynting's theorem is valid for any true orbit that solves the stiff equations of motion. Thinking of Poynting's theorem as detached from the complexity of the equations of motion might suggest a use of this theorem to calculate the radiation of hypothetical orbits that are not even solution to the equations of motion. For example, one can show that with half/retarded plus half/advanced fields a perfectly circular orbit does not radiate energy. But again, it is just as easy to show that a perfectly circular orbit is not a solution of the equations of motion with the self-interaction terms. Stiffness can prevent the dynamics from staying even for a single turn in the neighborhood of an orbit that would be a solution to the equation without the stiff terms [8]. A familiar use of the retarded-only radiation fields of DR is to calculate the incoherent dissipation of a circular current by applying Poynting's theorem to the radiation fields of a hypothetical circular orbit. It is a trivial exercise to show that this produces the same result of integrating the Lorentz-Dirac self-interaction times the velocity along that hypothetical circular orbit. Even though this approximation is fine for macroscopic currents restricted by a wire and forced by a battery, the stiff delay equations of electrodynamics can jump wildly given unstable conditions. Therefore, one should see first if the solution exists and if it is stable in the model and only then calculate the radiation for that precise stable orbit. The perturbations to circularity defined by Eq. (2) are necessary to stabilize the orbit, and as we show below, are essential as well to determine the radiated energy. The self-interaction force is a very small singular dissipative term along the unperturbed circular orbit, and the dissipation calculated over the perturbed orbit is completely changed by the gyroscopic perturbations, even if the final orbit stays near the circular orbit. After proper account

of these perturbations, our metastable orbit radiates even with the half-retarded plus half-advanced fields.

The half/retarded plus half/advanced radiation magnetic field of the electron in the DF model (the far-magnetic field) is [15]

$$B_{rad} = \frac{(a_- \times \hat{n}_-)}{2(1 - \hat{n}_- \cdot v_-)^2 r} - \frac{(a_+ \times \hat{n}_+)}{2(1 - \hat{n}_+ \cdot v_+)^2 r}, \quad (13)$$

where v and a are the electronic velocity and acceleration, $\hat{n}$ is a unit vector from the electron to the observation point, underscore minus indicates evaluation on the retarded light-cone and underscore plus indicates evaluation on the advanced light-cone. These two light-cones are defined by $t_{\pm} = t \pm (r - \hat{n}_{\pm} \cdot x)$ where x is the vector position of the electron measured from the center of the circular orbit. Once $(t_+ - t_-) \simeq 2r$, whenever $2r$ is an integer multiple of the distance travelled by light in a circular period (one light-period), the first term of expansion (13) along the precise circular orbit vanishes and the next term is the important one; a quadratic function of the orbital quantities. The same cancellation happens for the far-electric field whenever $2r$ is an odd-integer multiple of half the light-period, but for us here the far-magnetic field is enough. The next term of the magnetic expansion about the circular orbit is

$$B_{rad}^{(1)} = \frac{2(\hat{n} \cdot v)(a \times \hat{n}) + (\hat{n} \cdot x)(\dot{a} \times \hat{n})}{r}. \quad (14)$$

We can estimate (14) by noticing that along the $\hat{n}_{\pm} = \hat{x}$ direction of the unperturbed plane this quadratic term contains a product of the z perturbed coordinate times the x perturbed coordinate. Those perturbations are described by the u and Z nonrunaway normal modes already used and explained above Eq. (7). Using the normal mode conditions $\theta \dot{u} = \Omega \lambda_{xy} u$ and $\theta \dot{Z} = \Omega \lambda_z Z$ together with Eq.(8) one can show that the quadratic form uZ is a complex amplitude that oscillates harmonically with the beat frequency $(\lambda_{xy} + \lambda_z)\Omega/\theta = (\epsilon_1 - \epsilon_2)\Omega/\theta$. This complex amplitude is precisely the square-root of our resonant-normal form (7), which is a constant because it is the modulus of the complex number $uZ = \sqrt{C} \exp[i(\epsilon_1 - \epsilon_2)\Omega t/\theta]$. Translating the u mode to Cartesian coordinates with Eq. (2), we obtain the following approximation to $B_{rad}^{(1)}$

$$B_{rad}^{(1)} \propto \frac{2uZ}{r} \exp(i\Omega t), \quad (15)$$

and therefore the frequency of the emission line is equal to Ω plus the frequency of uZ

$$w_{DF} = \Omega + (\epsilon_1 - \epsilon_2)\Omega/\theta, \quad (16)$$

with Ω given by Eq. (1). Notice that the emitted frequency of the DF model is naturally different from the

orbital frequency. Among all possible settings of electrodynamics, only DF has this possibility. The fact that the emission frequency of hydrogen is different from the orbital frequency is another famous conundrum for the usual DR electromagnetic setting. The emission frequency of Eq. (16) contains differences of eigenvalues of the linear infinite-dimensional operator (5) and is strikingly similar to the Rydberg-Ritz combinatorial principle of quantum mechanics for the emission lines.

To discuss the width of the emission lines, we notice that for small amplitudes the resonant normal form constrains the deformation while allowing a radiative recoil. Even though the first term of the resonant normal form is a function of the relative separation only, a higher-order term might not be so, and at some point the recoil breaks the metastable orbit down. The construction of this next term shall be left for future work. In the dynamical process of resonant dissipation, the sharp line is emitted while the dynamics is locked in the neighborhood of the original orbit. We conjecture that when the metastable orbit breaks down, it can fall into the next metastable attractor; another circular orbit, or into the ground state. The stiff modes used in Eq. (8) describe a fast (stiff) time-scale of a frequency of the order of $\sigma/\theta \simeq 1400$ times the orbital frequency, the time for a stiff jump of the dynamics. After this fast timescale the resonance essentially locks the dynamics to the neighborhood of the metastable circular orbit.

The present simple approximation does not yet compete with the precision of QED, but it is surprising that recognizing the correct qualitative dynamical behavior of the equations of motion of classical electrodynamics can take us so far (the small corrections to Eq. (5) will be given elsewhere). The results are also beyond the Darwin approximation to Fokker's Lagrangian because the logarithm is not analytic at $\theta = 0$. We see then that the physics of resonant dissipation is qualitatively beyond the non-ionization criterion previously used by us[12], even though it contains it. Our stability condition forces the particles to stay bound during dissipation, a bound state condition analogous to the normalizability of the wave functions of Schroedinger's equation. Our stability analysis naturally involved an infinite-dimensional linear eigenvalue problem, Eq. (5), which is essentially a PDE, like Schroedinger's equation. We conjecture here that it should be possible to construct an average linear operator from Eq. (5) with $g = 1$ and Eq. (5) with $g = -7$ such that $\Omega + (\epsilon_1 - \epsilon_2)\Omega/\theta$ is a difference between its consecutive eigenvalues. That would be the Schroedinger's eigenvalue problem. Last, the angular momenta of Eq. (11) depend only logarithmically on the mass ratio times

θ^4 , such that the deuterium and the muonium atoms have essentially the same values for θ (i.e., the same fine structure constant, in agreement with QED). Notice that if the proton mass is set infinite, the quantized angular momenta become infinite as well, such that this logarithmic σ is a genuine two-body effect.

Recognizing the correct qualitative dynamical behavior is an intuition that goes a long way here; it suggested we should explore the infinite dimensionality of the delay with stability analysis and make use of the stiff modes. These stiff modes provide a natural integer number to label the metastable orbits. The natural appearance of the infinite-dimensional operator (Eq. (5)), and the large body of qualitative and quantitative agreement suggests that simple stability analysis inside the DF electrodynamics is a consistent new version of QED; one with an $\hbar$ given by Eq. (11). More work should be done to understand the relation between those two theories.

We thank Savio B. Rodrigues for discussions and for checking the numerical work in the complex λ plane with MATLAB.

* author's email address: deluca@df.ufscar.br

- [1] Staruszkiewicz A, 1968 *Acta Physica Polonica* **XXXIII**, 1007, see eq. (15).
- [2] Bohr N 1913 *Philos. Mag.* **26**, 1; Bohr N 1913 *Philos. Mag.* **26**, 476 .
- [3] Schild A 1963 *Phys. Rev.* **131**, 2762.
- [4] Andersen CM and Von Baeyer HC 1972 *Phys. Rev. D* **5**, 802.
- [5] Dirac PAM 1938 *Proc. R. Soc. London, Ser. A* **167**, 148 .
- [6] Currie DG, Jordan TF and Sudarshan ECG 1963 *Rev. Mod. Phys.* **35** 350, Marmo G, Mukunda N and Sudarshan ECG 1984 *Phys. Rev. D* **30** 2110
- [7] Driver RD 1979 *Phys. Rev. D* **19**, 1098 .
- [8] J. Grasman, *Asymptotic Methods for Relaxation Oscillations and Applications*, Applied Mathematical Sciences, **63**, Springer-Verlag, New York (1987).
- [9] Eliezer CJ 1943 *Proc. Cambridge Philos. Soc.* **39**, 173.
- [10] De Luca J 1998 *Phys. Rev. Lett.* **80**, 680 .
- [11] De Luca J 1998 *Phys. Rev. E* **58**, , 5727.
- [12] De Luca J 2000 *Phys. Rev. E* **62**, 2060.
- [13] Wheeler JA and Feynman RP 1945 *Rev. Mod. Phys.* **17** 157 ; Wheeler JA and Feynman RP 1949 **21** 425
- [14] R. E. Bellman and K.L.Cooke, *Differential-Difference Equations*, Academic Press, New York (1963), page 393.
- [15] F. Rohrlich, *Classical Charged Particles*, Addison-Wesley, New York (1965).
- [16] De Luca J, to be published.